

IDENTICAL PARTICLES AND PERMUTATION GROUP

*E.Celeghini*¹, *M.Rasetti*² and *G.Vitiello*³.

¹Dipartimento di Fisica - Università di Firenze and INFN–Firenze, I 50125 Firenze, Italy

²Dipartimento di Fisica and Unità INFN – Politecnico di Torino, I 10129 Torino, Italy

³Dipartimento di Fisica - Università di Salerno and INFN–Napoli, I 84100 Salerno, Italy

Abstract. Second quantization is revisited and creation and annihilation operators are shown to be related, on the same footing both to the algebra $h(1)$, and to the superalgebra $osp(1|2)$ that are shown to be both compatible with Bose and Fermi statistics. The two algebras are completely equivalent in the one-mode sector but, because of grading of $osp(1|2)$, differ in the many-particle case. The same scheme is straightforwardly extended to the quantum case $h_q(1)$ and $osp_q(1|2)$.

Claiming that a permutation of two particles has been performed requires distinguishability of the particles themselves. An idealized operational procedure to this effect would be for example the following: one first attaches a label to each particle (*i.e.* a quantum number identifying its state) in order to distinguish it from any other, then one interchanges the particles and, finally, one looks once more at the labels, to make sure that the exchange has been properly performed. However, one of the fundamental hypotheses of quantum field theory is exactly that particles should be treated as identical and indistinguishable. For this reason, the permutation group cannot be related *ab initio* to second quantization but, as we shall show, is to be introduced in the theory only at a second stage, by the physical request of expressing the properties on n -particle states in terms of first quantization observables.

This has deep consequences to the effect that the usual connection between algebraic properties of second quantization operators and statistics of the particles turns out to bear some arbitrariness. In order to prove this statement, we shall build explicitly, in terms of anticommuting creation and annihilation operators, a theory

where, by imposing the symmetry or antisymmetry of the particle states, both bosons and fermions can be simultaneously constructed. As briefly discussed at the end of the paper, the construction presented should be considered as an example of a much more general and far-reaching feature: since there is no necessary connection between the observables over the Fock space and the particle statistics, we are allowed not only to consider fermions and bosons as related to the Weyl-Heisenberg algebra $h(1)$, but the scheme is extendable also to more complex relation among observables (*e.g.* quantum algebras) and/or exotic statistics (*e.g.* anyons). It should be mentioned that the approach presented in this letter was inspired by the property that most of the algebraic structures relevant to second quantization physics are Hopf algebras^[1]: it is just the operation of coproduct, characteristic of the latter but usually disregarded because primitive for Lie algebras, recently brought to attention by studies of quantum algebras, which stands at the basis of our conceptual construction.

More formally, let us begin by showing how creation and annihilation operators can be related, on the same footing both to the algebra $h(1)$, and to the superalgebra $osp(1|2)$. $h(1)$, is customarily defined^[2] to be generated by the four operators $(a, a^\dagger, \mathbb{I}, N)$, with commutation relations

$$[a, a^\dagger] = \mathbb{I}, \quad [N, a] = -a, \quad [N, a^\dagger] = a^\dagger, \quad [\mathbb{I}, \bullet] = 0. \quad (1)$$

Upon characterizing the unitary representations (*i.e.* those for which $N^\dagger = N$, $(a^\dagger)^\dagger = a$) with lower bounded spectrum of N by the lowest eigenvalue n_0 , one can write

$$\begin{aligned} a^\dagger |k + n_0\rangle &= \sqrt{k+1} |k + n_0 + 1\rangle, & a |k + n_0\rangle &= \sqrt{k} |k + n_0 - 1\rangle, \\ N |k + n_0\rangle &= (k + n_0) |k + n_0\rangle, & k &\in \mathbb{N}. \end{aligned}$$

The usual Fock space $\mathcal{F}$ is obtained for $n_0 = 0$, usually adopting the relation $N \equiv a^\dagger a$ (one of the solutions of the equations $[N, a] = -a$, $[N, a^\dagger] = a^\dagger$):

$$\begin{aligned} a^\dagger |n\rangle &= \sqrt{n+1} |n+1\rangle, & a |n\rangle &= \sqrt{n} |n-1\rangle, \\ N |n\rangle &= n |n\rangle, & n &\in \mathbb{N}. \end{aligned} \quad (2)$$

A related $\mathbb{Z}_2$ -graded structure will be considered here, starting from the set of three operators $\mathcal{S} \equiv (a, a^\dagger, H)$ (notice that, as in (1), all operators are assumed to be unrelated), with H even and a and $a^\dagger$ odd. $\mathcal{S}$ is characterized by the relations

$$\{a, a^\dagger\} = 2H, \quad [H, a] = -a, \quad [H, a^\dagger] = a^\dagger, \quad (3)$$

and it is a *subset*, not a *sub-algebra*, of the $\mathbb{Z}_2$ -graded algebra $osp(1|2)$ [3]. Completion of $\mathcal{S}$ to the whole $osp(1|2)$ in fact requires the introduction of the additional set $\mathcal{S}' \equiv (J_-, J_+)$ in the even sector,

$$J_+ = \tfrac{1}{2}\{a^\dagger, a^\dagger\} \quad , \quad J_- = \tfrac{1}{2}\{a, a\} \quad , \quad (4)$$

with commutation relations

$$\begin{aligned} [J_+, a] &= -2a^\dagger \quad , & [J_-, a^\dagger] &= 2a \quad , & [J_+, a^\dagger] &= 0 = [J_-, a] \quad , \\ [J_+, J_-] &= -4H \quad , & [H, J_\pm] &= \pm 2J_\pm \quad . \end{aligned} \quad (5)$$

The bosonic sector $\mathcal{B} \equiv (J_-, J_+, \tfrac{1}{2}H)$ is, as well known, isomorphic to $su(1, 1)$, in the direct sum of the representations $\kappa = \frac{1}{4}, \frac{3}{4}$ [4].

An explicit analysis shows that the set $\mathcal{S}$ with relations (3) is sufficient to give rise to unitary representations of $osp(1|2)$ that have the spectrum of H bounded below, and can be characterized by the lowest eigenvalue of H , say h_0 . Explicitly,

$$\begin{aligned} a^\dagger |h_0 + 2k + 1\rangle &= \sqrt{2(k+1)} |h_0 + 2k + 2\rangle \quad , \\ a^\dagger |h_0 + 2k\rangle &= \sqrt{2(k+h_0)} |h_0 + 2k + 1\rangle \quad , \\ a |h_0 + 2k + 1\rangle &= \sqrt{2(k+h_0)} |h_0 + 2k\rangle \quad , \\ a |h_0 + 2k\rangle &= \sqrt{2k} |h_0 + 2k - 1\rangle \quad , \\ H |h_0 + k\rangle &= (h_0 + k) |h_0 + k\rangle \quad , \quad k \in \mathbb{N} \quad , \end{aligned} \quad (6)$$

where the partition of states in two classes exhibits the existence of supersymmetric doublets.

The main point of our derivation is the fact that eqs.(6), with $h_0 = \frac{1}{2}$, read

$$\begin{aligned} a^\dagger |h\rangle &= \sqrt{h + \tfrac{1}{2}} |h + 1\rangle \quad , \\ a |h\rangle &= \sqrt{h - \tfrac{1}{2}} |h - 1\rangle \quad , \\ H |h\rangle &= h |h\rangle \quad , \quad h \in \mathbb{N} + \tfrac{1}{2} \quad , \end{aligned} \quad (7)$$

namely they coincide with eqs.(2), provided the identification $h = n + \frac{1}{2}$ is implemented. This means that the closed subset $\mathcal{S}$ of $osp(1|2)$ defined by (3) shares the representation (2) – which can by induction be extended to the whole $osp(1|2)$ – with the Weyl-Heisenberg algebra $h(1)$. In other words, the Fock space $\mathcal{F}$ provides a faithful representation for both $h(1)$ (for $n_0 = 0$) and $osp(1|2)$ (for $h_0 = \frac{1}{2}$).

Second quantization is based, essentially, on the relations (2). We suggest that creation and annihilation operators may therefore be, with equal rights, be interpreted as belonging either to $osp(1|2)$ or to $h(1)$. The key point in our argument is

the following: as far as one considers the algebra as generated by the defining commutation relations only, any physical interpretation is contained in eqs. (2), and it turns out to be essentially irrelevant whether one selects $osp(1|2)$ or $h(1)$. However, when one deals with many-particle states, the two schemes lead to self-consistent yet mutually unequivalent descriptions. The reason why this may happen, is that $h(1)$ and $osp(1|2)$ are Hopf algebras (more precisely, $osp(1|2)$ is a *super* Hopf algebra). Any Hopf algebra, say $\mathcal{A}$, has, among its defining operations, the coproduct $\Delta : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (in fact, in the representation considered here, this is necessary, in that it implies that the action of the algebra is well defined on $\mathcal{F} \otimes \mathcal{F}$ and, by induction, on $\mathcal{F}^{\otimes n}$). For both $h(1)$ and $osp(1|2)$ Δ is, of course, primitive.

In $h(1)$ one has:

$$\begin{aligned} \Delta(a) &= a \otimes \mathbf{1} + \mathbf{1} \otimes a \equiv a_1 + a_2 \quad , \quad \Delta(a^\dagger) = a^\dagger \otimes \mathbf{1} + \mathbf{1} \otimes a^\dagger \equiv a_1^\dagger + a_2^\dagger \quad , \\ \Delta(N) &= N \otimes \mathbf{1} + \mathbf{1} \otimes N \equiv N_1 + N_2 \quad , \quad \Delta(\mathbb{I}) = \mathbb{I} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbb{I} \quad , \end{aligned} \quad (8)$$

(notice that $\mathbb{I}$ is a central operator with eigenvalue 1 and not the identity $\mathbf{1}$ of the universal enveloping algebra (UEA)).

The coalgebra for the superalgebra $osp(1|2)$ looks quite similar:

$$\begin{aligned} \Delta(a) &= a \otimes \mathbf{1} + \mathbf{1} \otimes a \equiv a_1 + a_2 \quad , \quad \Delta(a^\dagger) = a^\dagger \otimes \mathbf{1} + \mathbf{1} \otimes a^\dagger \equiv a_1^\dagger + a_2^\dagger \quad , \\ \Delta(H) &= H \otimes \mathbf{1} + \mathbf{1} \otimes H \equiv H_1 + H_2 \quad , \end{aligned} \quad (9)$$

however, since a and $a^\dagger$ are odd, whereas H is even, we have for $c, d, e, f \in osp(1|2)$, the multiplication law on $\mathcal{F} \otimes \mathcal{F}$

$$(c \otimes d)(e \otimes f) = (-)^{p(d)p(e)} ce \otimes df \quad , \quad (10)$$

where $p(d)$ and $p(e) \in \mathbb{Z}_2$ are the degrees (*i.e.* parities) of d and e respectively. On $\mathcal{F}^{\otimes n}$ composition rules are therefore quite different. Let us denote

$$\begin{aligned} a_j &\equiv \mathbf{1} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes a \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \quad , \\ a_j^\dagger &\equiv \mathbf{1} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes a^\dagger \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \quad , \\ H_j &\equiv \mathbf{1} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes H \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \quad , \\ N_j &\equiv \mathbf{1} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes N \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} \quad , \end{aligned}$$

where the multiple $\otimes$ -products have n factors, in each of which the only element different from the UEA identity $\mathbf{1}$ is in the j -th position. One has, for $(a, a^\dagger, N, \mathbb{I})$ in $h(1)$ the customary relations $[a_i, a_j] = 0$, $[a_i, a_j^\dagger] = \delta_{ij} \mathbb{I}$, $[N_i, a_j] = -a_i \delta_{ij}$ (plus their

hermitian conjugates), while for $(a, a^\dagger, H)$ in $osp(1|2)$, the (graded) commutation relations are

$$\begin{aligned}\{a_i, a_j^\dagger\} &= 2H_i \delta_{ij}, \quad , \quad [H_i, a_j] = -a_i \delta_{ij}, \quad , \quad [H_i, a_j^\dagger] = a_i^\dagger \delta_{ij} \quad , \\ \{a_i^\dagger, a_j^\dagger\} &= \delta_{ij} J_+ \quad , \quad \{a_i, a_j\} = \delta_{ij} J_- \quad ,\end{aligned}$$

and, of course, $[a_j, a_j] = 0$.

On the Fock basis of $\mathcal{F}^{\otimes n}$ adoption of $osp(1|2)$ leads to

$$\begin{aligned}a_j^\dagger |n_1, \dots, n_{j-1}, n_j, \dots, n_n\rangle &= (-1)^{s_j} \sqrt{n_j + 1} |n_1, \dots, n_{j-1}, n_j + 1, \dots, n_n\rangle \quad , \\ a_j |n_1, \dots, n_{j-1}, n_j, \dots, n_n\rangle &= (-1)^{s_j} \sqrt{n_j} |n_1, \dots, n_{j-1}, n_j - 1, \dots, n_n\rangle \quad , \\ N_j |n_1, \dots, n_{j-1}, n_j, \dots, n_n\rangle &= n_j |n_1, \dots, n_{j-1}, n_j, \dots, n_n\rangle \quad ,\end{aligned}\tag{11}$$

where the phases turn out to be exactly those customarily used in textbooks for

fermions ^[5] $\left(s_j \equiv \sum_{k=1}^{j-1} n_k\right)$.

It is worth pointing out that eqs.(11) differ from the usual (bosonic) ones only in the choice of phases and, on $\mathcal{F}^{\otimes n}$, imply $[a_i, a_i^\dagger] = \mathbb{I}$, consistently with the standard formulation, which in turn gives $\{a_i, a_i^\dagger\} = 2H_i (\equiv 2N_i + 1)$. Nevertheless the subtle and important implication here is that, in order to determine the phases of the basis vectors, an order must be imposed *a priori* in the set of indices j 's, in such a way that

$$|n_1, \dots, n_j, \dots, n_n\rangle \equiv \frac{1}{\sqrt{\prod_{j=1}^n n_j!}} (a_1^\dagger)^{n_1} \dots (a_j^\dagger)^{n_j} \dots (a_n^\dagger)^{n_n} |0\rangle \quad ,$$

contrary to the standard bosonic theory, where the creation operators commute and can be applied in any order. Of course, the usual properties of the Fock space, such as completeness

$$\sum_{\{n\}} |n_1, n_2, \dots\rangle \langle n_1, n_2, \dots| = \mathbb{I} \quad ,$$

and the projection operators on the one- and two-particles states

$$\mathcal{P}_1 \equiv \sum_i |1_i\rangle \langle 1_i| \quad , \quad \mathcal{P}_2 \equiv \sum_{i < j} |1_i, 1_j\rangle \langle 1_i, 1_j| + \sum_i |2_i\rangle \langle 2_i| \quad ,\tag{12}$$

where

$$\begin{aligned}|1_i\rangle &\equiv |n_1 = 0, n_2 = 0, \dots, n_{i-1} = 0, n_i = 1, n_{i+1} = 0, \dots, n_n = 0\rangle \quad , \\ |2_i\rangle &\equiv |n_1 = 0, n_2 = 0, \dots, n_{i-1} = 0, n_i = 2, n_{i+1} = 0, \dots, n_n = 0\rangle \quad , \\ |1_i, 1_j\rangle &\equiv |n_1 = 0, n_2 = 0, \dots, n_i = 1, \dots, n_j = 1, \dots, n_n = 0\rangle \quad ,\end{aligned}$$

do not depend on such phases, and persist. $osp(1|2)$ can, in such a way, be utilized to construct n -particle states, leading to a scheme non-equivalent to that derivable from $h(1)$, because of the grading of *odd* operators.

This possibility of using anticommuting operators, without restriction on the occupation numbers n_j 's, also casts a new light on the question of how one should introduce statistics into play.

As stressed in the introduction, second quantization is essentially uncorrelated with statistics^[6], which is required as a necessary set of rules to represent isomorphically n -particle states in the state space given by the n -fold tensorization of the single-particle Hilbert space, proper of first quantization. The relevant point in the analysis of this problem performed by Pauli in [6] is that the symmetry with respect to permutation of two particles does not depend on the prescription to build $\mathcal{F}$ from the vacuum (*i.e.* on the commutation or anticommutation relations of the a_i 's and $a_j^\dagger$'s), but it must be imposed as an external constraint aiming to guarantee a correct implementation of the above isomorphism.

In such a perspective, let us consider what happens with two bosons. Independently on whether the algebra is graded or not, one has to consider for such a system a symmetric Hilbert space, namely a generic state vector must be symmetric with respect to the exchange of the two particles (this being the feature which qualifies them as boson)

$$|x_1, x_2 \rangle_B \equiv \frac{1}{\sqrt{2}} (|x_1 \rangle |x_2 \rangle + |x_2 \rangle |x_1 \rangle) \quad ,$$

and, by (12),

$$|x_1, x_2 \rangle_B = \sum_{i < j} |1_i, 1_j \rangle \langle 1_i, 1_j | x_1, x_2 \rangle_B + \sum_i |2_i \rangle \langle 2_i | x_1, x_2 \rangle_B \quad .$$

Independently on how the states $|1_i, 1_j \rangle$ and $|2_i \rangle$ are constructed from the vacuum, the symmetry is here automatically implemented, in that

$$\begin{aligned} \langle 1_i, 1_j | x_1, x_2 \rangle_B &= \frac{1}{\sqrt{2}} \langle 1_i, 1_j | (|x_1, x_2 \rangle + |x_2, x_1 \rangle) \\ &= \frac{1}{\sqrt{2}} (\langle 1_i | x_1 \rangle \langle 1_j | x_2 \rangle + \langle 1_i | x_2 \rangle \langle 1_j | x_1 \rangle) \quad , \\ \langle 2_i | x_1, x_2 \rangle_B &= \frac{1}{\sqrt{2}} \langle 2_i | (|x_1, x_2 \rangle + |x_2, x_1 \rangle) = \langle 1_i | x_1 \rangle \langle 1_i | x_2 \rangle \quad , \end{aligned} \tag{13}$$

manifestly have the required invariance with respect to interchange of the two particles.

The feature that statistics has no connection with the algebra is further proved by the fact that $osp(1|2)$, as well as $h(1)$, work equally well with fermions. For two fermions we must consider an antisymmetric Hilbert state-space

$$|x_1, x_2 \rangle_F \equiv \frac{1}{\sqrt{2}} (|x_1 \rangle |x_2 \rangle - |x_2 \rangle |x_1 \rangle) \quad ,$$

and, once more independently on the algebra considered, the antisymmetry in the exchange of the two fermions is guaranteed, as well as the Pauli exclusion principle:

$$\begin{aligned}
\langle 1_i, 1_j | x_1, x_2 \rangle_F &= \frac{1}{\sqrt{2}} \langle 1_i, 1_j | (|x_1, x_2\rangle - |x_2, x_1\rangle) \\
&= \frac{1}{\sqrt{2}} (\langle 1_i | x_1 \rangle \langle 1_j | x_2 \rangle - \langle 1_i | x_2 \rangle \langle 1_j | x_1 \rangle) \quad , \\
\langle 2_i | x_1, x_2 \rangle_F &= \frac{1}{\sqrt{2}} \langle 2_i | (|x_1, x_2\rangle - |x_2, x_1\rangle) \\
&= \frac{1}{\sqrt{2}} (\langle 1_i | x_1 \rangle \langle 1_i | x_2 \rangle - \langle 1_i | x_2 \rangle \langle 1_i | x_1 \rangle) = 0 \quad .
\end{aligned} \tag{14}$$

(13) and (14) clearly demonstrate the novel possibility – besides the customary scheme^[4] – of constructing bosons with graded operators or fermions with even operators.

It should be stressed that our arguments in this paper are quite different from other procedures whereby "ad hoc" constraints are introduced on the variables in order to generate the statistics. An instance of such different approaches are non-linear transformations, along the lines proposed by Gutzwiller's projection operator method^[7]. In the fermionic case, a suggestive example is provided by the new creation and annihilation operators defined by

$$c_j = a_j \mathcal{Q}_j \quad , \quad \mathcal{Q}_j \equiv \frac{(\mathbf{1} - e^{i\pi N_j})}{\sqrt{N_j + \frac{1}{2}(\mathbf{1} + e^{i\pi N_j})}} = \mathcal{Q}_j^\dagger \quad , \quad c_j^\dagger = \mathcal{Q}_j a_j^\dagger \quad . \tag{15}$$

It is straightforward to check that – because of (2) – (15) leads to both the Pauli exclusion principle, $c_j^2 = 0 = (c_j^\dagger)^2$, and the customary fermionic anticommutation relations $\{c_i, c_j^\dagger\} = \delta_{ij} \mathbb{I}$, on the subsector of $\mathcal{F}^{\otimes n}$ consisting of paired superdoublets $\{|\dots, 2n_j, \dots\rangle, |\dots, 2n_j + 1, \dots\rangle\}$. The usual fermions can therefore be recovered by restriction to $n_j = 0$. Of course, in the above procedure no reference or use has been done of grading.

Adoption of a graded algebra has also deep bearings on the structures one can induce in the UEA . For example, it is usually assumed that the algebra $su(1, 1)$ can be constructed in the UEA of $h(1)$. This is not true: as stressed before $su(1, 1)$ can be easily obtained from eqs.(3) as the bosonic sector of the superalgebra $osp(1|2)$, while one has to use $n_0 = 0$ (*i.e.* one needs to impose indirectly the $osp(1|2)$ properties also) to obtain the same result from eqs.(1). Moreover, the grading property (10) plays an essential rôle in obtaining the coalgebra (of course primitive) of $su(1, 1)$ from the one of $\mathcal{S}$, while it is impossible to obtain the same result from the coalgebra of $h(1)$.

Indeed, from (5), (8) and (9), we have, for instance, in $h(1)$ $\Delta(J_-) = a^2 \otimes \mathbf{1} + \mathbf{1} \otimes a^2 + 2a \otimes a$ whereas in $osp(1|2)$, because of (11), $\Delta(J_-) = a^2 \otimes \mathbf{1} + \mathbf{1} \otimes a^2 \equiv J_- \otimes \mathbf{1} + \mathbf{1} \otimes J_-$, as it should, because J_- is primitive.

This shows that $su(1,1)$ is contained as a full Hopf algebra in the universal envelope of $\mathcal{S}$, while only in the common representation (2), the $su(1,1)$ algebra can be considered as realized in the UEA of $h(1)$ (of course, all these can be extended to $su(2)$ by analytical continuation from $su(1,1)$).

We have now to recall that both $h(1)$ ^[8] and $osp(1|2)$ ^[9] have quantum deformations and the whole discussion could be easily extended to them. Actually, it should be kept in mind that, in the scheme proposed, no relation links the algebraic features of the creation and annihilation operators to the symmetry of the states. It is, indeed, possible to study systems of particles, both fermions or bosons, by means of either $h_q(1)$ or $osp_q(1|2)$. The Fock space remain the same, while differences appear principally in the relations of composed observables with the single-particle ones.

We finally conjecture that in the present approach, there is room for considering objects with more complex symmetry such as anyons.

Acknowledgements

The authors gratefully acknowledge fruitful discussions with F. Iachello.

References.

- [1] J. Fuchs, *Affine Lie Algebras and Quantum Groups*, Cambridge University Press, Cambridge, 1992
- [2] W. Miller, *Lie Theory and Special Functions*, Academic Press, New York, NY, 1968
- [3] V.G. Kac, Comm. Math. Phys. **53**, 31 (1977)
P. Ramond, Physica **D 15**, 25 (1985)
- [4] A. Perelomov, *Generalized Coherent States and Their Applications*, Springer Verlag, Berlin, 1986
- [5] S.S.Schweber, *An Introduction to Relativistic Quantum Field Theory*, Row Peterson, Evanston ILL, 1961
- [6] W. Pauli, *Handbuch der Physik* **24** 1.Teil, 83 (1933) (english translation *General Principles of Quantum Mechanics*, Springer, Berlin, 1980)
- [7] M. Gutzwiller, Phys. Rev. **A 137**, 1726 (1965)
- [8] E. Celeghini, R. Giachetti, E. Sorace, and M. Tarlini, J. Math. Phys. **31**, 2548 (1990), and **32**, 1155 (1991)
- [9] P.P. Kulish, and N.Yu. Reshetikin, Lett. Math. Phys. **18**, 143 (1989)
E.Celeghini, T.D.Palev and M.Tarlini, Mod. Phys. Lett **B5**, 187 (1991)